




# Gulf of Mexico Harmful Algal Bloom Bulletin

Monday, January 08, 2018  
 NOAA National Ocean Service  
 NOAA Satellite and Information Service  
 NOAA National Weather Service

## Region: Southwest Florida



Instructions for viewing this geospatial pdf are available at: <https://go.usa.gov/xn9g2>.



## Conditions Report

Not present to medium concentrations of *Karenia brevis* (commonly known as Florida red tide) are present along- and offshore portions of southwest Florida, and not present in the Florida Keys. *K. brevis* concentrations are patchy in nature and levels of respiratory irritation will vary locally based upon nearby bloom concentrations, ocean currents, and wind speed and direction.

Health information, from the Florida Department of Health and other agencies, is available at [https://tidesandcurrents.noaa.gov/hab/gomx\\_health.html](https://tidesandcurrents.noaa.gov/hab/gomx_health.html). For recent, local observations and data check Mote Marine Laboratory Daily Beach Conditions (<http://visitbeaches.org>) and the Florida Fish and Wildlife Conservation Commission Red Tide Status (<http://myfwc.com/redtidestatus>).

### Recently Reported Impacts (Listed by County):

**Respiratory irritation:** Sarasota, Charlotte, and Lee  
**Dead fish:** Sarasota, Charlotte, and Lee  
**Discolored water:** None

### Definition of respiratory irritation levels.

| RESPIRATORY IRRITATION LEVEL | AFFECTED POPULATION |                               |                       |                                |                                   |
|------------------------------|---------------------|-------------------------------|-----------------------|--------------------------------|-----------------------------------|
|                              | NONE                | CHRONIC RESPIRATORY CONDITION | SENSITIVE TO RED TIDE | GENERAL PUBLIC (MILD SYMPTOMS) | GENERAL PUBLIC (INTENSE SYMPTOMS) |
| None                         | X                   |                               |                       |                                |                                   |
| Very low                     |                     | X                             |                       |                                |                                   |
| Low                          |                     | X                             | X                     |                                |                                   |
| Moderate                     |                     | X                             | X                     | X                              |                                   |
| High                         |                     | X                             | X                     | X                              | X                                 |

In the map above, the highest level of potential respiratory irritation forecast is displayed as a layer for each day from 01-08-18 to 01-11-18. See next page for a table of the respiratory irritation forecasts.


| State Name | County Region  | Mon 01/08 | Tue 01/09 | Wed 01/10 | Thu 01/11 |  |  |  |
|------------|--|-----------|-----------|-----------|-----------|--|--|--|
|            | DIXIE County-Gulf Coast                              |           |           |           |           |  |  |  |
|            | LEVY County-Gulf Coast                               |           |           |           |           |  |  |  |
|            | CITRUS County-Gulf Coast                             |           |           |           |           |  |  |  |
|            | HERNANDO County-Gulf Coast                           |           |           |           |           |  |  |  |
|            | Northern PASCO County-Gulf Coast                     |           |           |           |           |  |  |  |
|            | Southern PASCO County-Gulf Coast                     |           |           |           |           |  |  |  |
|            | Northern PINELLAS County-Gulf Coast                  |           |           |           |           |  |  |  |
|            | Northern PINELLAS County-Bay Regions                 |           |           |           |           |  |  |  |
|            | Northern PINELLAS County, Upper Bay Area-Bay Regions |           |           |           |           |  |  |  |
|            | Southern PINELLAS County-Gulf Coast                  | none      | none      | none      | none      |  |  |  |
|            | Southern PINELLAS County-Bay Regions                 |           |           |           |           |  |  |  |
|            | PINELLAS and Northern MANATEE County-Bay Regions     |           |           |           |           |  |  |  |
|            | South MANATEE County-Gulf Coast                      | very low  | low       | very low  | low       |  |  |  |
|            | South MANATEE County-Bay Regions                     | very low  | very low  | very low  | very low  |  |  |  |
|            | North SARASOTA County-Gulf Coast                     | very low  | low       | very low  | moderate  |  |  |  |
|            | North SARASOTA County-Bay Regions                    | very low  | very low  | very low  | very low  |  |  |  |
|            | Southern SARASOTA County-Gulf Coast                  | very low  | low       | very low  | moderate  |  |  |  |
|            | Southern SARASOTA County-Bay Regions                 | very low  | very low  | very low  | very low  |  |  |  |
|            | North CHARLOTTE County-Gulf Coast                    | very low  | low       | very low  | moderate  |  |  |  |
|            | North CHARLOTTE County-Bay Regions                   | very low  | very low  | very low  | very low  |  |  |  |
|            | Southern CHARLOTTE County-Gulf Coast                 | very low  | low       | very low  | moderate  |  |  |  |
|            | Southern CHARLOTTE County-Bay Regions                | moderate  | moderate  | moderate  | moderate  |  |  |  |
|            | Upper CHARLOTTE Harbor-Bay Regions                   |           |           |           |           |  |  |  |
|            | Northern LEE County-Gulf Coast                       | very low  | low       | very low  | low       |  |  |  |
|            | Northern LEE County-Bay Regions                      | low       | very low  | low       | low       |  |  |  |
|            | Central LEE County-Gulf Coast                        | very low  | low       | very low  | low       |  |  |  |

The table lists the highest level of potential respiratory irritation forecast. *K. brevis* concentrations are patchy in nature and levels of respiratory irritation will vary locally based upon nearby bloom concentrations, ocean currents, and wind speed and direction.


| State Name | County Region                       | Mon 01/08 | Tue 01/09 | Wed 01/10 | Thu 01/11 |  |  |  |
|------------|-------------------------------------|-----------|-----------|-----------|-----------|--|--|--|
|            |                                     |           |           |           |           |  |  |  |
|            | Central LEE County-Bay Regions      | low       | very low  | low       | low       |  |  |  |
|            | Southern LEE County-Gulf Coast      |           |           |           |           |  |  |  |
|            | Southern LEE County-Bay Regions     |           |           |           |           |  |  |  |
|            | Northern COLLIER County-Gulf Coast  |           |           |           |           |  |  |  |
|            | Northern COLLIER County-Bay Regions |           |           |           |           |  |  |  |
|            | Central COLLIER County-Gulf Coast   |           |           |           |           |  |  |  |
|            | Central COLLIER County-Bay Regions  |           |           |           |           |  |  |  |
|            | Southern COLLIER County-Gulf Coast  |           |           |           |           |  |  |  |
|            | Northern MONROE County-Gulf Coast   |           |           |           |           |  |  |  |
|            | Southern MONROE County-Gulf Coast   |           |           |           |           |  |  |  |
|            | UPPER KEYS-Oceanside                |           |           |           |           |  |  |  |
|            | UPPER KEYS and FLORIDA BAY-Gulfside |           |           |           |           |  |  |  |
|            | MIDDLE KEYS-Oceanside               |           |           |           |           |  |  |  |
|            | MIDDLE KEYS-Gulfside                |           |           |           |           |  |  |  |
|            | LOWER KEYS-Oceanside                |           |           |           |           |  |  |  |
|            | LOWER KEYS-Gulfside                 |           |           |           |           |  |  |  |

The table lists the highest level of potential respiratory irritation forecast. *K. brevis* concentrations are patchy in nature and levels of respiratory irritation will vary locally based upon nearby bloom concentrations, ocean currents, and wind speed and direction.

### Wind conditions from Venice Pier, FL



Wind speed and direction are averaged over 12 hours from buoy measurements. Length of line indicates speed; angle indicates direction. Red indicates that the wind direction favors upwelling near the coast. Values to the left of the dotted vertical line are measured values; values to the right are forecasts. Wind observation and forecast data provided by NOAA's National Weather Service (NWS). A text summary of the marine forecast by region is available from NWS at <https://go.usa.gov/xnx4y>.



## Analysis

### Summary of Recent Water Samples:

#### ***K. brevis* Cell Concentrations:**

**Range:** Not Present through Medium

**Date:** 12/29-01/04

**Source:** FWRI, MML, SCHD, CCPCD

### Imagery:

Recent ensemble imagery (MODIS Aqua, 1/7) is partially obscured by clouds from Pinellas to Charlotte counties, limiting analysis. Patches of elevated to very high chlorophyll (2 to >20 µg/L) with the optical characteristics of *K. brevis* are present along- and offshore southwest Florida from Charlotte to Monroe counties.


### Forecasts:

Observed winds over the weekend may have promoted southerly transport of surface bloom concentrations.

Davis, Ludema

*Karenia brevis* cell concentration sampling data from: 12/29/17 through 01/04/18. Cell count data are provided by Florida FWC Fish and Wildlife Research Institute. For a list of sample providers and a key to the cell concentration categories, please see the HAB-OFS bulletin guide: [https://tidesandcurrents.noaa.gov/hab/hab\\_publication/GOMX\\_HAB\\_Bulletin\\_Guide.pdf](https://tidesandcurrents.noaa.gov/hab/hab_publication/GOMX_HAB_Bulletin_Guide.pdf). Detailed sample information can be obtained through the Florida FWC Fish and Wildlife Research Institute: <http://myfwc.com/REDTIDESTATUS>.

MODIS satellite chlorophyll image (01/07/18) with possible *K. brevis* HAB areas shown by red polygon(s).



*Karenia brevis* cell concentration sampling data from: 12/29/17 through 01/04/18. Cell count data are provided by Florida FWC Fish and Wildlife Research Institute. For a list of sample providers and a key to the cell concentration categories, please see the HAB-OFS bulletin guide: [https://tidesandcurrents.noaa.gov/hab/hab\\_publication/GOMX\\_HAB\\_Bulletin\\_Guide.pdf](https://tidesandcurrents.noaa.gov/hab/hab_publication/GOMX_HAB_Bulletin_Guide.pdf). Detailed sample information can be obtained through the Florida FWC Fish and Wildlife Research Institute: <http://myfwc.com/REDTIDESTATUS>. MODIS satellite chlorophyll image (01/07/18).

Verified and suspected HAB areas shown in red. Other areas with *K. brevis* optical characteristics shown in yellow (see p. 4 analysis for interpretation).