



# Gulf of Mexico Harmful Algal Bloom Bulletin

Monday, June 25, 2018

NOAA National Ocean Service


NOAA Satellite and Information Service

NOAA National Weather Service

## Region: Southwest Florida



Instructions for viewing this geospatial pdf are available at: <https://go.usa.gov/xn9g2>.



In the map above, the highest level of potential respiratory irritation forecast is displayed as a layer for each day from 06-25-18 to 06-28-18. See next page for a table of the respiratory irritation forecasts.

## Conditions Report

Not present to high concentrations of *Karenia brevis* (commonly known as red tide) are present along- and offshore portions of southwest Florida, and not present in the Florida Keys. *K. brevis* concentrations are patchy in nature and levels of respiratory irritation will vary locally based upon nearby bloom concentrations, ocean currents, and wind speed and direction.

### Recently Reported Impacts (Listed by County):

**Respiratory irritation:** Sarasota, Lee, Collier

**Dead fish:** Sarasota, Charlotte, Collier

**Discolored water:** Sarasota

### Definition of respiratory irritation levels.

RESPIRATORY IRRITATION LEVEL	AFFECTED POPULATION				
	NONE	CHRONIC RESPIRATORY CONDITION	SENSITIVE TO RED TIDE	GENERAL PUBLIC (MILD SYMPTOMS)	GENERAL PUBLIC (INTENSE SYMPTOMS)
None	X				
Very low		X			
Low		X	X		
Moderate		X	X	X	
High		X	X	X	X

## Additional Resources

### Health Information:

#### Florida Department of Health:

<http://www.floridahealth.gov/environmental-health/aquatic-toxins/red-tide.html>

**Other resources:** <https://go.usa.gov/xQNWp>

### Recent, Local Observations and Data:

#### Mote Marine Laboratory Daily Beach Conditions:

<http://visitbeaches.org>

#### Florida Fish and Wildlife Conservation Commission:

<http://myfwc.com/redtidestatus>

State Name	County Region	Mon 06/25	Tue 06/26	Wed 06/27	Thu 06/28			
	DIXIE County-Gulf Coast							
	LEVY County-Gulf Coast							
	CITRUS County-Gulf Coast							
	HERNANDO County-Gulf Coast							
	Northern PASCO County-Gulf Coast							
	Southern PASCO County-Gulf Coast							
	Northern PINELLAS County-Gulf Coast							
	Northern PINELLAS County-Bay Regions							
	Northern PINELLAS County, Upper Bay Area-Bay Regions							
	Southern PINELLAS County-Gulf Coast	none	none	none	none			
	Southern PINELLAS County-Bay Regions							
	PINELLAS and Northern MANATEE County-Bay Regions	none	none	none	none			
	South MANATEE County-Gulf Coast	none	none	none	none			
	South MANATEE County-Bay Regions	low	low	low	low			
	North SARASOTA County-Gulf Coast	high	high	high	high			
	North SARASOTA County-Bay Regions	moderate	moderate	moderate	moderate			
	Southern SARASOTA County-Gulf Coast	high	high	high	high			
	Southern SARASOTA County-Bay Regions							
	North CHARLOTTE County-Gulf Coast	moderate	moderate	moderate	moderate			
	North CHARLOTTE County-Bay Regions							
	Southern CHARLOTTE County-Gulf Coast	moderate	moderate	moderate	moderate			
	Southern CHARLOTTE County-Bay Regions	moderate	moderate	moderate	moderate			
	Upper CHARLOTTE Harbor-Bay Regions							
	Northern LEE County-Gulf Coast	moderate	high	high	high			
	Northern LEE County-Bay Regions	none	none	none	none			
	Central LEE County-Gulf Coast							

The table lists the highest level of potential respiratory irritation forecast. *K. brevis* concentrations are patchy in nature and levels of respiratory irritation will vary locally based upon nearby bloom concentrations, ocean currents, and wind speed and direction.


Cells are marked 'none' if *K. brevis* was detected, but no respiratory irritation is forecasted in the region. Cells are blank if no *K. brevis* has been detected in the region.

State Name	County Region	Mon 06/25	Tue 06/26	Wed 06/27	Thu 06/28			
	Central LEE County-Bay Regions							
	Southern LEE County-Gulf Coast							
	Southern LEE County-Bay Regions							
	Northern COLLIER County-Gulf Coast							
	Northern COLLIER County-Bay Regions							
	Central COLLIER County-Gulf Coast	very low	low	low	very low			
	Central COLLIER County-Bay Regions	low	low	low	low			
	Southern COLLIER County-Gulf Coast							
	Northern MONROE County-Gulf Coast							
	Southern MONROE County-Gulf Coast							
	UPPER KEYS-Oceanside							
	UPPER KEYS and FLORIDA BAY-Gulfside							
	MIDDLE KEYS-Oceanside							
	MIDDLE KEYS-Gulfside							
	LOWER KEYS-Oceanside							
	LOWER KEYS-Gulfside							

The table lists the highest level of potential respiratory irritation forecast. *K. brevis* concentrations are patchy in nature and levels of respiratory irritation will vary locally based upon nearby bloom concentrations, ocean currents, and wind speed and direction.


Cells are marked 'none' if *K. brevis* was detected, but no respiratory irritation is forecasted in the region. Cells are blank if no *K. brevis* has been detected in the region.


### Wind conditions from Naples, FL



Wind speed and direction are averaged over 12 hours from buoy measurements. Length of line indicates speed; angle indicates direction. Red indicates that the wind direction favors upwelling near the coast. Values to the left of the dotted vertical line are measured values; values to the right are forecasts. Wind observation and forecast data provided by NOAA's National Weather Service (NWS). A text summary of the marine forecast by region is available from NWS at <https://go.usa.gov/xnx4y>.

### Wind conditions from Venice Pier, FL





## Analysis

### Summary of Recent Water Samples:

#### ***K. brevis* Cell Concentrations:**

**Range:** Not Present through High

**Date:** 06/15-06/21

**Source:** (FWRI, MML, SCHD, CCPCD)

### Imagery:

Recent ensemble imagery (MODIS Aqua, 6/23) is mostly obscured south of Charlotte County, limiting analysis. A patch of elevated to very high chlorophyll (3 to >20 µg/L) with the optical characteristics of *K. brevis* is present alongshore Sarasota and Charlotte counties.


### Forecasts:

Forecasted winds today through Thursday may increase the potential for onshore transport of surface *K. brevis* concentrations and the potential for respiratory irritation.

Ludema, Yang

*Karenia brevis* cell concentration sampling data from: 06/15/18 through 06/21/18. Cell count data are provided by Florida FWC Fish and Wildlife Research Institute. For a list of sample providers and a key to the cell concentration categories, please see the HAB-OFS bulletin guide: [https://tidesandcurrents.noaa.gov/hab/hab\\_publication/GOMX\\_HAB\\_Bulletin\\_Guide.pdf](https://tidesandcurrents.noaa.gov/hab/hab_publication/GOMX_HAB_Bulletin_Guide.pdf). Detailed sample information can be obtained through the Florida FWC Fish and Wildlife Research Institute: <http://myfwc.com/REDTIDESTATUS>.

MODIS Aqua satellite chlorophyll image (06/23/18) with possible *K. brevis* HAB areas shown by red polygon(s).



*Karenia brevis* cell concentration sampling data from: 06/15/18 through 06/21/18. Cell count data are provided by Florida FWC Fish and Wildlife Research Institute. For a list of sample providers and a key to the cell concentration categories, please see the HAB-OFS bulletin guide: [https://tidesandcurrents.noaa.gov/hab/hab\\_publication/GOMX\\_HAB\\_Bulletin\\_Guide.pdf](https://tidesandcurrents.noaa.gov/hab/hab_publication/GOMX_HAB_Bulletin_Guide.pdf). Detailed sample information can be obtained through the Florida FWC Fish and Wildlife Research Institute: <http://myfwc.com/REDTIDESTATUS>. MODIS Aqua satellite chlorophyll image (06/23/18).

Verified and suspected HAB areas shown in red. Other areas with *K. brevis* optical characteristics shown in yellow (see p. 4 analysis for interpretation).