



# Gulf of Mexico Harmful Algal Bloom Bulletin


Monday, January 11, 2021

NOAA National Ocean Service

NOAA Satellite and Information Service

NOAA National Weather Service

Instructions for viewing this geospatial pdf are available at: <https://go.usa.gov/xn9g2>.



The image above is the top layer in a series of maps for 01-11-21 to 01-14-21 displaying the highest level of potential respiratory irritation forecasts in each region.

## Region: Southwest Florida



### Conditions Report

Not present to high concentrations of *Karenia brevis* (commonly known as red tide) are present along- and offshore portions of southwest Florida, and not present in the Florida Keys. *K. brevis* concentrations are patchy in nature and levels of respiratory irritation will vary locally based upon nearby bloom concentrations, ocean currents, and wind speed and direction.

Recently Reported Impacts (Listed by County):

**Respiratory irritation:** Lee, Collier

**Dead fish:** Lee, Collier

Definition of respiratory irritation levels.

RESPIRATORY IRRITATION LEVEL	AFFECTED POPULATION				
	NONE	CHRONIC RESPIRATORY CONDITION	SENSITIVE TO RED TIDE	GENERAL PUBLIC (MILD SYMPTOMS)	GENERAL PUBLIC (INTENSE SYMPTOMS)
None	X				
Very low		X			
Low		X	X		
Moderate		X	X	X	
High		X	X	X	X

### Additional Resources

Health Information:

**Florida Department of Health:**

<http://www.floridahealth.gov/environmental-health/aquatic-toxins/harmful-algae-blooms/index.html>

**Other resources:** <https://go.usa.gov/xQNWp>

Recent, Local Observations and Data:

**Mote Marine Laboratory Daily Beach Conditions:**

<http://visitbeaches.org>

**Florida Fish and Wildlife Conservation Commission:**

<http://myfwc.com/redtidestatus>

State Name	County Region	Mon 01/11	Tue 01/12	Wed 01/13	Thu 01/14			
Florida								
	DIXIE County-Gulf Coast							
	LEVY County-Gulf Coast							
	CITRUS County-Gulf Coast							
	HERNANDO County-Gulf Coast							
	Northern PASCO County-Gulf Coast							
	Southern PASCO County-Gulf Coast							
	Northern PINELLAS County-Gulf Coast							
	Northern PINELLAS County-Bay Regions							
	Northern PINELLAS County, Upper Bay Area-Bay Regions							
	Southern PINELLAS County-Gulf Coast							
	Southern PINELLAS County-Bay Regions							
	PINELLAS and Northern MANATEE County-Bay Regions							
	South MANATEE County-Gulf Coast							
	South MANATEE County-Bay Regions							
	North SARASOTA County-Gulf Coast							
	North SARASOTA County-Bay Regions							
	Southern SARASOTA County-Gulf Coast							
	Southern SARASOTA County-Bay Regions							
	North CHARLOTTE County-Gulf Coast							
	North CHARLOTTE County-Bay Regions							
	Southern CHARLOTTE County-Gulf Coast							
	Southern CHARLOTTE County-Bay Regions	none	very low	very low	very low			
	Upper CHARLOTTE Harbor-Bay Regions							
	Northern LEE County-Gulf Coast	very low	low	very low	very low			
	Northern LEE County-Bay Regions	moderate	moderate	moderate	moderate			
	Central LEE County-Gulf Coast	moderate	moderate	low	low			

The table lists the highest level of potential respiratory irritation forecast. *K. brevis* concentrations are patchy in nature and levels of respiratory irritation will vary locally based upon nearby bloom concentrations, ocean currents, and wind speed and direction.


Cells are marked 'none' if *K. brevis* was detected, but no respiratory irritation is forecasted in the region. Cells are blank if no *K. brevis* has been detected in the region.

State Name	County Region	Mon 01/11	Tue 01/12	Wed 01/13	Thu 01/14			
Florida								
	Central LEE County-Bay Regions	moderate	moderate	moderate	moderate			
	Southern LEE County-Gulf Coast	moderate	moderate	low	low			
	Southern LEE County-Bay Regions	moderate	moderate	moderate	moderate			
	Northern COLLIER County-Gulf Coast	moderate	moderate	low	low			
	Northern COLLIER County-Bay Regions							
	Central COLLIER County-Gulf Coast	moderate	moderate	low	low			
	Central COLLIER County-Bay Regions	moderate	moderate	moderate	moderate			
	Southern COLLIER County-Gulf Coast							
	Northern MONROE County-Gulf Coast							
	Southern MONROE County-Gulf Coast							
	UPPER KEYS-Oceanside							
	UPPER KEYS and FLORIDA BAY-Gulfside							
	MIDDLE KEYS-Oceanside							
	MIDDLE KEYS-Gulfside							
	LOWER KEYS-Oceanside							
	LOWER KEYS-Gulfside							

The table lists the highest level of potential respiratory irritation forecast. *K. brevis* concentrations are patchy in nature and levels of respiratory irritation will vary locally based upon nearby bloom concentrations, ocean currents, and wind speed and direction.


Cells are marked 'none' if *K. brevis* was detected, but no respiratory irritation is forecasted in the region. Cells are blank if no *K. brevis* has been detected in the region.


**Wind conditions from Naples, FL**



Wind speed and direction are averaged over 12 hours from buoy measurements. Length of line indicates speed; angle indicates direction. Red indicates that the wind direction favors upwelling near the coast. Values to the left of the vertical line are measured values; values to the right are forecasts. Wind observation and forecast data provided by NOAA's National Weather Service (NWS). A text summary of the marine forecast by region is available from NWS at <https://www.weather.gov/marine/stheastmz>.

**Wind conditions from Venice Pier, FL**





## Analysis

### Summary of Recent Water Samples:

#### ***K. brevis* Cell Concentrations:**

**Range:** Not Present through High

**Date:** 01/01-01/07

**Source:** FWRI, MML, SCHD, CCPCD

### Imagery:

Recent satellite imagery has been obscured by clouds alongshore southwest Florida, limiting analysis. Imagery from 1/9 (VIIRS), continues to show very high chlorophyll (2 to >20 µg/L) is present alongshore southern Lee and Collier counties. Due to cloud cover, the region of chlorophyll containing the optical characteristics of *K. brevis* is not clearly defined; though recent beach reports indicate the bloom extent is unchanged from last week.

### Forecasts:

Variable to alongshore winds forecast today through Thursday (1/11-14) will promote southward transport of surface *K. brevis* concentrations and minimize any significant potential for respiratory irritation at the coast.


Additional satellite imagery available here:

<https://tidesandcurrents.noaa.gov/hab/gomx/data/Imagery-EasternGOMX/>


-Davis, Keeney

*Karenia brevis* cell concentration sampling data from: 01/01/21 through 01/07/21. Cell count data are provided by Florida FWC Fish and Wildlife Research Institute. For a list of sample providers and a key to the cell concentration categories, please see the HAB-OFS bulletin guide: [https://tidesandcurrents.noaa.gov/hab/hab\\_publication/GOMX\\_HAB\\_Bulletin\\_Guide.pdf](https://tidesandcurrents.noaa.gov/hab/hab_publication/GOMX_HAB_Bulletin_Guide.pdf). Detailed sample information can be obtained through the Florida FWC Fish and Wildlife Research Institute: <http://myfwc.com/REDTIDESTATUS>.

VIIRS satellite chlorophyll image (01/09/21) with possible *K. brevis* HAB areas shown by red polygon(s).



*Karenia brevis* cell concentration sampling data from: 01/01/21 through 01/07/21. Cell count data are provided by Florida FWC Fish and Wildlife Research Institute. For a list of sample providers and a key to the cell concentration categories, please see the HAB-OFS bulletin guide: [https://tidesandcurrents.noaa.gov/hab/hab\\_publication/GOMX\\_HAB\\_Bulletin\\_Guide.pdf](https://tidesandcurrents.noaa.gov/hab/hab_publication/GOMX_HAB_Bulletin_Guide.pdf). Detailed sample information can be obtained through the Florida FWC Fish and Wildlife Research Institute: <http://myfwc.com/REDTIDESTATUS>. VIIRS satellite chlorophyll image (01/09/21).



Verified and suspected HAB areas shown in red. Other areas with *K. brevis* optical characteristics shown in yellow (see p. 4 analysis for interpretation).